California

PROVIDER*Update*

NEWS & ANNOUNCEMENTS

SEPTEMBER 30, 2019

| UPDATE 19-736

Help Patients Better Control Their Asthma and Its Triggers

Together we can educate members to better manage their asthma

To support your work with patients with asthma, Health Net* is providing members who have asthma with educational materials about asthma, asthma medication compliance and managing their asthma triggers.

Why target asthma?

Asthma is one of the most prevalent, expensive health conditions in the U.S. In California alone, over 2 million adults and over 850,000 children currently have asthma.¹ When patients manage their asthma well, it can help them avoid emergency room visits and inpatient stays.²

Use these best practices for successful asthma care

According to the U.S. Department of Health and Human Services (HHS), successful asthma care must include proper diagnosis and treatment, followed by long-term regular follow-up to ensure control.³

- Asthma self-management education is essential to reducing asthma-related adverse health effects. Education improves quality of life by reducing urgent care visits, emergency room visits, hospitalizations, and health care costs.⁴
- The key to successfully managing asthma is ensuring patients know when to use controller medications and when rescue medications are needed to handle emergency asthma attacks.
- Consider using peak flow or spirometry with a physical exam to confirm a diagnosis of asthma.
- Once you establish a confirmed asthma diagnosis, manage your patients' use of controller and rescue medications.
 - If patients use more rescue medication than controller medication, it may indicate that a patient's current medication regimen is not working and needs reassessment. Identify patient barriers and work with the patient to ensure their controller medication(s) are managing their asthma well.

Improve asthma control with these helpful tips and tools

- Create an asthma action plan. A written action plan will help patients recognize early warning signs, symptoms and peak expiratory flow (PEF) measures that indicate worsening asthma.
- Make sure patients have an asthma spacer. Make sure your patients with asthma have a spacer. When patients use a spacer, they will get more medication

Health Net® 3 PAGES

THIS UPDATE APPLIES TO CAL MEDICONNECT PROVIDERS:

- Physicians
- Participating Physician Groups
- Hospitals
- Ancillary Providers

PROVIDER SERVICES

provider_services@healthnet.com Los Angeles County – 1-855-464-3571 San Diego County – 1-855-464-3572 www.healthnet.com

PROVIDER COMMUNICATIONS provider.communications@ healthnet.com

^{*} Health Net Community Solutions, Inc. is a subsidiary of Health Net, LLC and Centene Corporation. Health Net is a registered service mark of Health Net, LLC. All other identified trademarks/service marks remain the property of their respective companies. All rights reserved. CONFIDENTIALITY NOTE FOR FAX TRANSMISSION: This facsimile may contain confidential information. The information is intended only for the use of the individual or entity named above. If you are not the intended recipient, or the person responsible for delivering it to the intended recipient, you are hereby notified that any disclosure, copying, distribution, or use of the information contained in this transmission is strictly PROHIBITED. If you have received this transmission in error, please notify the sender immediately by telephone or by return fax and destroy this transmission, along with any attachments.

into their lungs and less medication deposited in the mouth and throat. It also helps them coordinate inhaling and pressing on their metered dose inhaler.

• Teach patients about a peak flow meter (PFM). Make sure patients have a peak flow meter and know how to use it.


To check the specific health plan coverage for spacers and PFMs for your patient(s), please review their *Schedule of Benefits* or contact Provider Services at the telephone number as listed in the right-hand column on page 1.

Provide pre-discharge review. Review your patient's inhaler technique and, when possible, environmental control
strategies to reduce the patient's exposure to allergens and irritants.

Improve health outcomes with culturally sensitive and linguistically appropriate care

- Speak with your patients about their cultural views and beliefs about asthma. Address your patient's concerns, beliefs and attitudes, as they are key to successful asthma self-management.
- **Provide asthma education that is age, culturally and linguistically appropriate.** Once you have identified your patient's cultural and language needs, be sure to discuss the following:
 - Confirm your patients and their families and/or support persons understand the:
 - Asthma diagnosis and basic facts about asthma.
 - Severity of their asthma.
 - Patient's particular asthma triggers and how to minimize exposure to them.
- Initiate a prescription medication, if needed. Explain how the medications work and how some are required to be used even though your patient may not be having symptoms.
- Teach the patient how to use an inhaler and spacer. Demonstrate their use.
- · Complete a written asthma action plan. Review the plan with the patient.
- Schedule a follow-up appointment. This will help you see how well the patient is able to control their asthma and if they are following your medication recommendations.

Refer to the national asthma care guidelines at www.nhlbi.nih.gov/files/docs/guidelines/asthma_qrg.pdf, and strive to deliver culturally and linguistically appropriate care at every recommended phase of each visit. For a quick look at each phase, see the Asthma Quick Reference⁵ figure below.



Use correct codes for conditions other than asthma

To help ensure that members with asthma are accurately identified, be sure to also code for those members with the respiratory conditions listed below, if applicable.

Codes for Conditions Other than Asthma

Description	Codes	
Emphysema	J43.0, J43.1, J43.2, J43.8, J43.9	
Other emphysema	J98.2–J98.3	
COPD	J44.0, J44.1, J44.9	
Chronic respiratory conditions due to fumes/vapors	J68.4	
Cystic fibrosis	E84.0, E84.11, E84.19, E84.8–E84.9	
Acute respiratory failure	J96.00–J96.02, J96.20–J96.22	

Additional information

Providers are encouraged to access Health Net's provider portal online at provider.healthnet.com for real-time information, including eligibility verification, claims status, prior authorization status, plan summaries, and more.

If you have questions regarding the information contained in this update, contact the Health Net Provider Services Center by county within 60 days at:

Line of Business	Telephone Number	Email Address
Cal MediConnect – Los Angeles County	1-855-464-3571	<pre>provider_services@healthnet.com</pre>
Cal MediConnect – San Diego County	1-855-464-3572	

¹ https://www.cdph.ca.gov/Programs/CCDPHP/DEODC/EHIB/CPE/CDPH%20Document%20Library/Asthma_Surveillance_in_CA_Report_2017.pdf.

² www.ncqa.org/hedis/measures/medication-management-for-people-with-asthma-and-asthma-medication-ratio/.

³ U.S. Department of Health and Human Services. National Institute of Health. National Heart, Lung, and Blood Institute. Asthma Care Quick Reference. Diagnosing and Managing Asthma. September 2012. Accessed online August 2016 at www.nhlbi.nih.gov/health-pro/guidelines/current/asthma-guidelines/quick-reference.

⁴ www.cdc.gov/asthma/pdfs/asthma_facts_program_grantees.pdf.

⁵ https://www.nhlbi.nih.gov/files/docs/guidelines/asthma_qrg.pdf. U.S. Department of Health and Human Services. National Institute of Health. National Heart, Lung, and Blood Institute. Asthma Care Quick Reference. Diagnosing and Managing Asthma. September 2012. Accessed online August 2016 at www.nhlbi.nih.gov/health-pro/guidelines/current/asthma-guidelines/quick-reference.