

## Learn How to Improve Your HEDIS<sup>1</sup> Rate for Osteoporosis Screening

Use this tip sheet to review key details of the Osteoporosis Screening in Older Women (OSW) measure, best practices and resources.

Measure	Women 65–75 years of age who received osteoporosis screening.		
	<ul> <li>One or more osteoporosis screening tests on or between the member's 65th birthday and December 31 of the measurement year.</li> </ul>		
Exclusions	Patients who meet the following criteria:		
	<ul> <li>Had a claim/encounter for osteoporosis therapy any time in the member's history through December 31 of the year prior to the measurement year.</li> </ul>	Often called a silent disease, osteoporosis reduces bone mass through structural deterioration, resulting in compromised bone strength. Osteoporosis is more common in women than in men. One in two women will have an osteoporosis-related fracture in their lifetime. <sup>2</sup> Routine screenings can help identify osteoporosis and allow members	
	• Received a dispensed prescription to treat osteoporosis anytime on or between January 1 three years prior to the measurement year through December 31 of the year prior to the measurement year.		
	<ul><li>Hospice or using hospice services anytime during the measurement year.</li><li>Palliative care during the measurement year.</li></ul>		
	<ul> <li>Members 66 and older as of December 31 of the measurement year enrolled in an Institutional Special Needs Plan (I-SNP) or living long-term in an institution any time during the measurement year.</li> </ul>	to take preventive measures.	
	• Members 66 and older as of December 31 of the measurement year with frailty and advanced illness.		

(continued)

Best practices	<ul> <li>Submit a claim for osteoporosis screening.</li> <li>Set up a process for bone mineral density (BMD) testing: <ul> <li>To determine if a member had a BMD test, query enrolled members in the age span noted.</li> <li>Provide a BMD prescription and urge patients to complete the screening soon</li> <li>Make sure the screening site notifies you with results in a timely manner.</li> <li>Set up a follow-up visit to discuss the results at the patient's next visit.</li> </ul> </li> <li>Evaluate women for risk factors that would increase the risk of osteoporosis. Some risk factors include low body weight, excessive alcohol intake, current tobacco use, history of fractures, and use of certain medications.</li> <li>Discuss osteoporosis prevention methods with your patients, such as calcium, vitamin D supplements, and weight-bearing exercises.</li> <li>Have culturally appropriate, tailored educational messaging that addresses the risk and importance of osteoporosis in racial and ethnic minority populations that have lower screening rates.</li> </ul>				
	Use the appropriate service codes when billing for OSW screenings CPT Copyright 2017 American Medical Association. All rights reserved. CPT® is a registered trademark of the American Medical Association.				
Codes	Tests/medications	CPT codes	HCPCS codes		
	Osteoporosis screening tests	76977, 77078, 77080, 77081, 77085			
	Osteoporosis medications		J0897, J1740, J3110, J3111, J3489		
	Long-acting osteoporosis medications during an inpatient stay		J0897, J1740, J3489		

## Resources

- Final Recommendation Statement, Osteoporosis to Prevent Fractures: Screening. U.S. Preventive Services Task Force at
- www.uspreventiveservicestaskforce.org/uspstf/recommendation/osteoporosis-screening.
- National Committee for Quality Assurance (NCQA), HEDIS Measurement Year 2022, Volume 2: Technical Specifications for Health Plans, 2021.

\*Health Net of California, Inc., Health Net Community Solutions, Inc. and Health Net Life Insurance Company are subsidiaries of Health Net, LLC and Centene Corporation. Health Net is a registered service mark of Health Net, LLC. All other identified trademarks/service marks remain the property of their respective companies. All rights reserved.