

## AUDIT-C - Overview

The AUDIT-C is a 3-item alcohol screen that can help identify persons who are hazardous drinkers or have active alcohol use disorders (including alcohol abuse or dependence). The AUDIT-C is a modified version of the 10 question AUDIT instrument.

### Clinical Utility

The AUDIT-C is a brief alcohol screen that reliably identifies patients who are hazardous drinkers or have active alcohol use disorders.

### Scoring

The AUDIT-C is scored on a scale of 0-12.

Each AUDIT-C question has 5 answer choices. Points allotted are:

a = 0 points, b = 1 point, c = 2 points, d = 3 points, e = 4 points

- **In men**, a score of 4 or more is considered positive, optimal for identifying hazardous drinking or active alcohol use disorders.
- **In women**, a score of 3 or more is considered positive (same as above).
- However, when the points are all from Question #1 alone (#2 & #3 are zero), it can be assumed that the patient is drinking below recommended limits and it is suggested that the provider review the patient's alcohol intake over the past few months to confirm accuracy.<sup>3</sup>
- Generally, the higher the score, the more likely it is that the patient's drinking is affecting his or her safety.

### Psychometric Properties

For identifying patients with heavy/hazardous drinking and/or Active-DSM alcohol abuse or dependence

	Men <sup>1</sup>	Women <sup>2</sup>
≥3	Sens: 0.95 / Spec. 0.60	Sens: 0.66 / Spec. 0.94
≥4	Sens: 0.86 / Spec. 0.72	Sens: 0.48 / Spec. 0.99

For identifying patients with active alcohol abuse or dependence

≥ 3	Sens: 0.90 / Spec. 0.45	Sens: 0.80 / Spec. 0.87
≥ 4	Sens: 0.79 / Spec. 0.56	Sens: 0.67 / Spec. 0.94

1. Bush K, Kivlahan DR, McDonnell MB, et al. The AUDIT Alcohol Consumption Questions (AUDIT-C): An effective brief screening test for problem drinking. *Arch Internal Med.* 1998 (3): 1789-1795.

2. Bradley KA, Bush KR, Epler AJ, et al. Two brief alcohol-screening tests from the Alcohol Use Disorders Identification Test (AUDIT): Validation in a female veterans affairs patient population. *Arch Internal Med Vol* 163, April 2003: 821-829.

3. Frequently Asked Questions guide to using the AUDIT-C can be found via the website: [www.oqp.med.va.gov/general/uploads/FAQ%20AUDIT-C](http://www.oqp.med.va.gov/general/uploads/FAQ%20AUDIT-C)

## AUDIT-C Questionnaire

Patient Name \_\_\_\_\_ Date of Visit \_\_\_\_\_

**1. How often do you have a drink containing alcohol?**

- ☐ a. Never
- ☐ b. Monthly or less
- ☐ c. 2-4 times a month
- ☐ d. 2-3 times a week
- ☐ e. 4 or more times a week

**2. How many standard drinks containing alcohol do you have on a typical day?**

- ☐ a. 1 or 2
- ☐ b. 3 or 4
- ☐ c. 5 or 6
- ☐ d. 7 to 9
- ☐ e. 10 or more

**3. How often do you have six or more drinks on one occasion?**

- ☐ a. Never
- ☐ b. Less than monthly
- ☐ c. Monthly
- ☐ d. Weekly
- ☐ e. Daily or almost daily